

Positive forward rates in the maximum smoothness framework

Julián Manzano*

Division of Applied Physics.

Institute of Physics and Measurement Technology.

Linköping University.

and

Jörgen Blomvall

Division of Optimization. Department of Mathematics.

Linköping University.

Abstract

In this article we present a non-linear dynamic programming algorithm for the computation of forward rates within the maximum smoothness framework. The algorithm implements the forward rate positivity constraint for a one-parametric family of smoothness measures and it handles price spreads in the constraining dataset. We investigate the outcome of the algorithm using the Swedish Bond market showing examples where the absence of the positive constraint leads to negative interest rates. Furthermore we investigate the predictive accuracy of the algorithm as we move along the family of smoothness measures. Among other things we observe that the inclusion of spreads not only improves the smoothness of forward curves but also significantly reduces the predictive error.

May 2003

*corresponding author: Julián Manzano, Division of Applied Physics. Institute of Physics and Measurement Technology. Linköping University, S-581 83 Linköping, Sweden; e-mail: manzano@ifm.liu.se; phone: +46.737.353.373.

1 Introduction

Within the financial industry forward rate curves play a central role in fixed-income derivative pricing and risk management. Notwithstanding, such curves are not empirical or directly measurable objects but rather useful abstract concepts from where observed prices can be derived. Moreover, given a finite set of market prices, we can construct in general an infinite number of compatible forward rate curves¹. To avoid such ambiguity several approaches have been proposed in the literature trying to capture a “reasonable” or “natural” functional form within the set of compatible possibilities.

In historical order the first kind of methods proposed to solve this problem make use of the so-called parametric approach. In this approach a particular functional form for the forward rate curve is assumed leaving a certain number of free parameters to be fixed from the calculation of a given set of quoted prices. An extensive literature exists advocating for this approach. We can cite as examples the works of McCulloch [1], Vasicek and Fong [2], Chambers, Carleton and Waldman [3], Shea [4], Nelson and Siegel [5] and more recently the works of Svensson [6], Fisher, Nychka and Zervos [7] and Waggoner [8]. In most of these works we notice the privileged role played by polynomial and exponential splines as the preferred functional forms for the forward rate curves.

The second kind of methods has been termed in the literature as non-parametric or maximum-smoothness approach. Here instead of advocating for an a priori functional form for the forward rate a given measure of smoothness is chosen and then the forward rate curve is obtained as the one maximizing this measure subject to the constraints imposed by market prices. Examples where these methods have been investigated include the works of Adams and Van Deventer [9], Delbaen and Lorimier [10], Kian Guan Lim, Qin Xiao et al [11, 12], Frishling and Yamamura [13] and Yekutieli [14]. In these works three different smoothing measures have been proposed. We also have the works of Forsgren [15] and Kwon [16] that generalize these methodologies and clarify the connection between splines and certain smoothness measures. Finally we point out the work of Wets, Bianchi and Yang [17] that can be located somewhere in-between both approaches since here the number of functional parameters is finite (albeit arbitrarily large) and the functional behavior is restricted to a subfamily of C^2 curves.

The purpose of this article is two-fold: Firstly we want to present an efficient maximum-smoothing algorithm that handles the presence of spreads and implements the positivity constraint. Secondly we want to investigate the predictive power of a linear combination of two quadratic measures, namely the one proposed by Delbaen et al. and Frishling et al.

¹We assume here that prices do not allow for arbitrage. If this is not the case the compatibility condition has to be relaxed.

[10, 13] and the one by Adams and van Deventer [9]. Here it is worth remarking that once the compatibility with market prices is fulfilled the only guiding principle that should be taken as definition of “reasonable” or “natural” is the predictive power and not other ad-hoc criteria.

In this article we will only use as constraining data coupon bearing bonds. The inclusion of treasury bills, zero coupon bonds or bill futures is straightforward and amounts to adding the corresponding linear constraints. Since our objective in this article is focusing on an algorithm dealing with non-linear constraints and inequality constraints (spreads and positivity constraint) we have not included such data.

With these objectives in mind we organize the article as follows:

In section 2 we present the objective function that we will use throughout the article and we establish the basic notation. In this section we present a sketch of the complete algorithm leaving the details for the appendices. In section 3 we present the results of the article including examples where the absence of the positivity constraint or the adequate spreads leads to negative rates. Here we present also a study of the predictive power of the one-parametric family of smoothness measures that include as extreme cases the measures used by Delbaen et al., Frishling et al. and Adams and Van Deventer. Finally in section 4 we present the conclusions.

2 The algorithm

A bond, j , is an instrument that gives future coupons, c_{ij} , at time stages $R_i^{(j)}$, $i = 1, \dots, n_j - 1$ and a final payment, $c_{n_j j} = N_j$ ². The bond price, P_j , can be determined from the discrete forward rate curve, f_r , $r = 1, \dots, R_{n_j}^{(j)}$, as follows

$$P_j = \sum_{i=1}^{n_j} c_{ij} \exp \left(- \sum_{r=1}^{R_i^{(j)}-1} f_r \xi_r \right), \quad (1)$$

where ξ_r is the length of the time period between time stage r and $r+1$ (in our implementation we have used $\xi_r = 1$ day).

The objective function, or smoothing measure³, is defined as a linear combination of the one used by Delbaen et al. and Frishling et al. (DF) [10, 13] and the one used by Adams and

²By final payment N_j we mean the complete last cash flow of bond j , typically that includes a principal plus a last coupon.

³Note that since we do not include a global minus sign we have to perform a minimization and not a maximization. With this sign, that is the one used in the literature, “rugosity” measure would be a more appropriate name.

$$W := \frac{\gamma}{2} \sum_{r=1}^{n-1} \left(\frac{f_{r+1} - f_r}{\xi_r} \right)^2 \xi_r + \frac{\varphi}{2} \sum_{r=2}^{n-1} \left(\frac{2}{(\xi_{r-1} + \xi_r)} \left(\frac{f_{r+1} - f_r}{\xi_r} - \frac{f_r - f_{r-1}}{\xi_{r-1}} \right) \right)^2 \xi_r, \quad (2)$$

The first term in Eq.(2) (DF) is a discrete approximation of the integral of the square of the first derivative of f and the second term (AD) is a discrete approximation of the integral of the square of the second derivative of f . This objective function is to be minimized subject to the consistency constraints

$$0 = \rho_j + \sum_{r=1}^{R_{n_j}^{(j)}-1} f_r \xi_r - \ln(v_j), \quad (3)$$

$$f_r \geq 0, \quad \rho_j^b \leq \rho_j \leq \rho_j^a, \quad (4)$$

where $\rho_j = \ln(P_j/N_j)$ is to be determined along with f and where we have used the definitions

$$\begin{aligned} v_j &:= \sum_{i=1}^{n_j} \alpha_{ij} \exp \left(\sum_{r=R_i^{(j)}}^{R_{n_j}^{(j)}-1} f_r \xi_r \right), & \rho_j^b &:= \ln(\alpha_{0j}^b), & \rho_j^a &:= \ln(\alpha_{0j}^a), \\ \alpha_{ij} &:= \frac{c_{ij}}{N_j}, & \alpha_{0j}^b &:= \frac{P_j^b}{N_j}, & \alpha_{0j}^a &:= \frac{P_j^a}{N_j} & \alpha_{n_j j} &:= 1, \end{aligned} \quad (5)$$

with P_j^b , P_j^a the respective bid and ask prices of bond j ($j = 1, \dots, m$)⁴. Note that the equality constraint given by Eq.(3) is just Eq.(1) rewritten taking logarithms and using the definitions (5). Eq.(4) introduces two inequality constraints. The first one is the positivity constraint over the forward rate curve and the second one is the requirement that the single price given by Eq.(1) must lie in-between the bid and ask prices.

We define the spread of bond j as the quantity $\rho_j^a - \rho_j^b$. We take the largest time to maturity in Eq.(2) equal to the largest time to maturity in the constraining dataset, namely

$$n := \max_j (R_{n_j}^{(j)}).$$

The constraints reflecting bond prices (1) have been rewritten in a way such that they become linear when no coupons are present ($v_j = 1$ for a zero-coupon bond j). Constraints given by Eqs.(3) and (4) are moved to the objective function defining

$$Z := W + \sum_{j=1}^m \lambda_j \left[\rho_j + \sum_{r=1}^{R_{n_j}^{(j)}-1} f_r \xi_r - \ln(v_j) \right] - \mu \sum_{r=1}^n \ln(f_r) - \tilde{\mu} \sum_{j=1}^m \left(\ln(\rho_j - \rho_j^b) + \ln(\rho_j^a - \rho_j) \right), \quad (6)$$

⁴Note that this optimization problem is non-convex and therefore several local minima may exist.

with the Lagrange multipliers λ_j , $j = 1, \dots, m$ and the logarithmic barriers with parameters $\mu > 0$ and $\tilde{\mu} > 0$ (in the solution procedure we take $\mu \rightarrow 0$, $\tilde{\mu} \rightarrow 0$). The use of log barriers to deal with inequality constraints is a standard methodology in interior point methods for optimization problems [18]. An explanation of this methodology adapted to our problem is given in appendix A. Briefly the minimization algorithm is structured as follows:

- step 0: Initialize log barriers with coefficients $\mu^{[0]}$, $\tilde{\mu}^{[0]}$ and set $(f, \rho) = (f^{[0]}, \rho^{[0]})$ with the seed $(f^{[0]}, \rho^{[0]})$ satisfying the inequality constraints (4). Let $k = 0$.
- step 1: Make a second order approximation of Z at $(f^{[k]}, \rho^{[k]})$ (see appendix A).
- step 2: Determine the newton step $(\hat{f}^{[k+1]}, \hat{\rho}^{[k+1]})$ using dynamic programming (see appendix B).
- step 3: Modify $(\hat{f}^{[k+1]}, \hat{\rho}^{[k+1]})$ to get a solution $(f^{[k+1]}, \rho^{[k+1]})$ that satisfies the inequality constraints (4). Update log barriers ($\mu^{[k]} \rightarrow 0$ and $\tilde{\mu}^{[k]} \rightarrow 0$ as $k \rightarrow \infty$) and calculate the values of W and of constraints (3). Check if a termination criterion is satisfied, otherwise let $k = k + 1$ and go to step 1 (see appendix C). (7)

Computing times involved in step 2 are summarized in subsection B.1. The solution typically stabilizes in approximately 6 iterations as can be seen in Fig.(1). On a Pentium 4, 2.4 Ghz computer the algorithm coded in C++ takes around 1/4 sec. to compute a forward rate curve like any of the ones seen in Fig.(2).

3 Consistency and predictability

In this section we present some examples of the behavior of algorithm (7) and we investigate the predictive power of the resulting forward rate curves. In Fig.(2) we present a series of forward rate curves calculated using DF and AD smoothing measures. There we can see that for both measures the resulting curves share some similar traits like the positions of most peaks and dips. Clear differences between both sets of curves are found at their end-points and in their behavior in the presence of high spreads. In the set of curves obtained from the DF measure we have vanishing first derivative at the end-points and curves that tend to constants for high spreads. For the AD measure we have vanishing second and third derivatives at end-points [16] and curves for high enough spreads given by straight lines. From the financial point of view these features are, in principle, just different aesthetic possibilities. In order to choose a particular measure the guiding principles should be, in the first place, the fulfillment of the consistency constraints given by Eqs.(3-4) and after this is guaranteed the predictive performance.

Let us start now with the analysis of consistency. As can be found in [16] if we do not




Figure 1: Plot of the objective function W (see Eq. (2)) vs. Newton iterations (starting from the seeds: $f^{[0]} = 0.04/\text{year}$, $\rho_j^{[0]} = (\rho_j^a + \rho_j^b)/2$). The plots correspond to the forward rate curves of Fig.(2). The converge behavior is affected by the termination criterion dictated by Eq.(29) and by the initial values of the log barriers dictated by Eq.(30). Note the exponential convergence of the algorithm in the first steps. Note also that on Friday 06 the minimal solution for AD measure is, for all practical purposes, a straight line.




Figure 2: Forward rate curves obtained from the prices of 11 Swedish government bonds in a two-week period of July 2001. The curves in plot (a) are calculated using DF measure ($\gamma = 1 \text{ year}^3, \varphi = 0$) while the ones in plot (b) correspond to the AD measure ($\gamma = 0, \varphi = 1 \text{ year}^5$). In both plots we have used a tolerance spread of 1% ($\rho_j^a - \rho_j^b = 0.01$).

consider the positivity constraint the local minima of objective (2) are given by exponential splines with exponents $\pm\sqrt{\gamma/\varphi}$ or polynomial splines of order 2 or 4 when φ or γ are respectively zero. The main problem with these exponential or polynomial splines is that there is no warranty that they fulfill the positivity constraint. Negative rates are not admissible in the absence of arbitrage opportunities and the risk of obtaining this unwanted feature is illustrated in Fig.(3). In this figure we have concentrated on the Swedish bond data on Monday, July 09, 2001. There we have tested three spread patterns for both DF and AD measures with and without the positivity constraint. From these plots it is evident that the inclusion of spreads in the calculation of the forward rates is a necessary ingredient that can have a major impact in the resulting functional behavior.

Once we have an algorithm that insures the consistency of forward rates we can concentrate on the predictive accuracy of different measures. However, before starting the analysis of this issue let us make a brief digression to comment a point regarding measure (2). If we want to have both γ and φ different from zero and we want to compare the effects of each term it is important to realize that DF and AD measures scale differently under changes of time units. In other words, γ and φ have different units. A practical way to define their units is to consider the objective W as an adimensional quantity. By doing so and remembering that f has units of inverse time, it is immediate to obtain that γ has units of time³ and φ units of time⁵. The importance of keeping this in mind becomes apparent in results like the ones presented in Figs.(4) and (5).

Figs.(4) and (5) summarize our results regarding the predictive accuracy of the algorithm as a function $\sqrt{\gamma/\varphi}$. There it is clear that the characteristic time span where DF and AD compete is not the day or the century, but clearly the year. To construct these figures we have calculated the forward rate curves for different values of $\sqrt{\gamma/\varphi}$ when one bond is removed from the constraining dataset. The price of this missing bond is used afterwards as a benchmark to test the accuracy of the resulting curves. Since we are interested in the statistical performance we have done such comparison for 335 consecutive trading days starting on Wednesday, November 08, 2000 and ending on Thursday, March 07, 2002.

We are also interested in studying the impact of spreads in the constraining dataset over the predictive accuracy. Therefore we present our results for three spread patterns, namely constant spreads of 0%, 0.5% and 1% in the constraining dataset. Fig.(4) concentrates on the predictive accuracy for the first 9 bonds of Table (2) and Fig.(5) presents the same analysis for the remaining 2 bonds of Table (2). These last 2 bonds are the ones with the larger maturities in the complete dataset. In particular for the last one with the largest maturity we have to decide upon the methodology to extrapolate the forward curve outside the range

of the constraining dataset⁵. Therefore in Fig.(5) we present the prediction accuracy using constant extrapolation from the last maturity in the constraining dataset and “ W -generated” extrapolation that consists in utilizing the W -optimal forward rate curve even outside the range of the constraining dataset.

4 Results and conclusions

In this article we have presented a non-linear dynamic programming algorithm designed for the calculation of positive definite forward rate curves using data with or without spreads. We have included multiple details of the algorithm aiming at practitioners not familiar with the techniques of dynamic programming. We have illustrated the results of this algorithms using the Swedish bond data for a one-parametric family of smoothness measures.

The results and conclusions are the following:

- The proposed algorithm calculates forward rate curves in real time and admits, without time or complexity penalizations, the use of any non-linear *local* objective function. Since it also handles non-linear constraints it is possible to include within the constraining dataset any derivative products with prices bearing some dependence on forward rates.
- This is the first algorithm proposed in the literature that implements the positivity constraint in the maximum smoothness framework. To the knowledge of the authors the only other work that implements such constraint outside this framework is the one of Wets et al. [17]. The proposal in [17] has the advantage of using simple linear programming but do not consider the presence of spreads minimizing instead the sum over the *modulus* of the difference between calculated prices and market prices.
- For the objective functions and constraining datasets like the ones we have used or more generally for the ones studied in [16] the algorithm proposed in that reference offers better computing times at the expense of ignoring the positivity constraint. Essentially the complexity in [16] is c^3 and in ours is $n c^2$ where n is the number of time steps and c the number of constraints. For that reason, when this class of objective functions and constraining datasets are used, a well coded algorithm might try first the proposal in [16] (improving its treatment of the spreads using e.g. log-barriers) and later, only if the result is not positive definite, use our approach.

⁵By the range of the constraining dataset we mean the range in time to maturity that goes from present to the last maturity within the dataset.




Figure 3: Forward rate curves obtained from the prices of 11 Swedish government bonds on Monday, July 09, 2001. The vertical lines indicate cash flows. Highest lines indicate the 11 cash flows at maturities (normalized to 1) and the smaller ones coupon cash flows. In these plots we show the effect of the positivity constraint with three spread patterns both for the DF (in (a1), (b1) and (c1)) and the AD (in (a2), (b2) and (c2)) measures. In plots (c1) and (c2) we see that a spread of 1% is enough to generate positive and sensible curves for both measures (the positivity constraint has no effect in this case). In plots (a1) and (a2) we take null spreads observing that for both measures large negative rates are obtained (conspicuously for AD measure). Also in these plots we see that even though the positivity constraint can be fulfilled the resulting curves have large oscillations. Negative rates or large oscillations in the positively constrained curves can be related to price patterns that are close to violate absence of arbitrage (the precise nature of this relation will be studied elsewhere). Plots (b1) and (b2) explores this fact keeping null spreads in two particular bonds (bonds SO 1043 and SO 1034 in Tables 1 and 2) and allowing for a large 5 % in the rest (almost unconstraining this subset). Doing so we observe in (b1) and (b2) curves that are still negative in the region where the nominal cash flows of the 0-spread bonds take place.




Figure 4: Plots indicating some statistical features of price prediction errors as a function of $\sqrt{\gamma/\varphi}$. These prediction errors are obtained removing the indicated bonds from the constraining dataset. The statistical sample comprise the forward curves corresponding to 335 consecutive trading days starting on Wednesday, November 08, 2000 and ending on Thursday, March 07, 2002. Within this set of days we calculate the relative errors of the predicted prices given by $(P^{pred.} - P^{market}) / P^{market}$ where P^{market} is the actual price of the bond given by the market and $P^{pred.}$ is the price predicted using the prices of all other bonds in the dataset. Hence in plots (a1), (a2) and (a3) we show the average of the absolute value of these errors and the plots (b1), (b2) and (b3) just their average. Consecutive columns show averages obtained using spreads of 0%, 0.5% and 1% in the constraining dataset. The set of predicted bonds shown in this figure is given by the first 9 bonds of Table 2 . The remaining bonds of this table (the last two bonds with the larger maturities) are analyzed in Fig.(5). Note how prediction errors tend to decrease when spreads are considered albeit not significantly (compare with Fig.(5)). Note also that in the absence of spreads the DF measure ($\sqrt{\gamma/\varphi} = \infty$) systematically exhibits a better performance than the AD one ($\sqrt{\gamma/\varphi} = 0$).




Figure 5: In this figure we consider the same kind of statistical features as in Fig.(4) but now the plots correspond to the last two bonds of Table 2 that are the ones with larger maturities within our dataset (we have also included the bond with the shortest maturity to facilitate the comparison with Fig.(4)). For the bond with the largest maturity the forward curve has to be extrapolated to reach its last cash flows (see Fig.(3)). We consider two extrapolation possibilities, one where we continue the forward rate curve from the last cash flow as a constant and the other where the curve is dictated by the minimization of functional W even beyond the last cash flow in the constraining dataset. Note that constant extrapolation gives better results than W-generated extrapolation in the region close to the AD measure ($\sqrt{\gamma/\phi} = 0$). Again like in Fig.(4) we observe that in the absence of spreads the DF measure ($\sqrt{\gamma/\phi} = \infty$) systematically exhibits a better performance than the AD one. The most striking difference with Fig.(4) is that here we observe that for these two long maturing bonds the introduction of spreads drastically reduce the prediction error all along the family of measures.

- Since the optimization problem we are solving is non-convex we can not discard the presence of several local minima (this is independent of the presence of the positivity constraint [16]). However, we have tested our algorithm starting from different seeds and in all cases we have arrived at the same minima. These tests included hundreds of searches with initial log-prices given by $\rho_j^{[0]} = \rho_j^b + x(\rho_j^a - \rho_j^b)$ (with $x \in [0, 1]$ a flat random variable) and initial forward rates given by several constant and oscillating functions. With this comment we only want to convey our practical experience and by no means we intend to say that we have exhaustively explored the presence or absence of local minima.
- It is clear that one way to avoid negative rates is just by increasing spreads by hand. The advantage of using real spreads at a given moment is that one can be sure of being consistent with market prices. If one observes in Fig.(3) the large forward rate variations taking place for different spread patterns no doubts should remain about the relevance of a careful treatment of this issue.
- From Fig.(5) we conclude that the inclusion of spreads can remarkably improve the accuracy of resulting forward curves in the prediction of market prices of long maturing bonds. In [16] it was pointed out that the inclusion of spreads notably improved the smoothness of the forward curve. To the knowledge of the authors this is the first time it is shown that their presence also improves the prediction accuracy. For that reason we believe that spreads should be considered even when the market data does not provide such information. In that case the approach should consist in using a cross validation technique to asses the optimal spread minimizing an error criterion based in the prediction of market prices (for cross validation methods see for example [20, 21]). One possibility is using the well know “leave-one-out” cross-validation to select the optimal spread much in the spirit suggested by Figs.(4) and (5). Given a constraining set of k products, the method actually consists in obtaining k forward rate curves for a given spread, each curve leaving out one of the constraining products (bonds in our case) but using only the k omitted products to compute an error criterion like $\sum_i^k \left| P_i^{pred.} - P_i^{market} \right| / P_i^{market}$. Thus we obtain a quantitative criterion to select an optimal level of spread⁶. For example for our dataset and our family of models (measures) it can be conjectured from Figs.(4) and (5) that this optimal level of spread is typically around 0.5%.

⁶This technique is stated here only to illustrate the existence of an optimal non vanishing level of spread whenever no real spreads are provided by the market. We do not intend to provide an efficient procedure to find this level.

- Figs.(4) and (5) strongly suggest that for low spread patterns DF measure is more accurate than AD one. For bigger spreads results do not clearly favor any particular measure.

Acknowledgements

J. M. acknowledges the financial support from *Tekniska Högskolan, Inst. för Fysik och Mätteknik*, Linköping University, thanks J. Shaw (Barclays) for calling his attention to ref. [17], F. Delbaen for ref. [10] and J. M. Eroles (HSBC) for proof reading the manuscript. J. M. would also like to thank Dr. Per Olov Lindberg for the hospitality during his stay at the Division of Optimization; Linköping University. The authors would like to thank the referees for their helpful comments on the manuscript.

Appendix A. Newton steps

Since the objective function Z in Eq.(6) is a non-quadratic function of f_r, ρ_j we will use an iterative quadratic approximation (Newton steps) to find its minima. We start with feasible seeds $f_r^{[0]}$ and $\rho_j^{[0]}$ fulfilling inequality constraints (4) and we set up initial log-barriers coefficients $\mu^{[0]} > 0$ and $\tilde{\mu}^{[0]} > 0$. For newton iteration number s we define

$$\begin{aligned}\hat{f}_r^{[s]} &:= f_r^{[s-1]} + \Delta_r^{[s]}, \\ \hat{\rho}_j^{[s]} &:= \rho_j^{[s-1]} + \sigma_j^{[s]},\end{aligned}\tag{8}$$

with $s = 1, 2, \dots$. The hat over f and ρ indicate that at each step s , $\hat{f}^{[s]}$ and $\hat{\rho}^{[s]}$ are the minima of the quadratic approximation and may not fulfill the inequality constraints (4). To assure constraints (4) are fulfilled a final redefinition $\hat{f}^{[s]} \rightarrow f^{[s]}, \hat{\rho}^{[s]} \rightarrow \rho^{[s]}$ is necessary after each Newton step. This redefinition is explained in appendix C. Expanding Z up to second order in $\Delta_r^{[s]}$ and $\sigma_j^{[s]}$ we write

$$\begin{aligned}Z^{[s]} &:= Z\left(\hat{f}^{[s]}, \hat{\rho}^{[s]}, \lambda^{[s]}\right)\Big|_{O(2)} \\ &= \frac{1}{2}\Delta^{[s]T}Q^{[s]}\Delta^{[s]} + \Delta^{[s]T}B^{[s]}\lambda^{[s]} + \Delta^{[s]T}C^{[s]} + \lambda^{[s]T}a^{[s]} \\ &\quad + \sigma^{[s]T}\lambda^{[s]} + \frac{1}{2}\sigma^{[s]T}M^{[s]}\sigma^{[s]} + \sigma^{[s]T}D^{[s]} + b^{[s]},\end{aligned}\tag{9}$$

where $b^{[s]}$ collects all terms not depending on $\Delta^{[s]}$, $\sigma^{[s]}$ or $\lambda^{[s]}$. We will use square brackets around Newton step indices and parenthesis around dynamic programming ones. Let us now work-out the matrices involved in the quadratic approximation. From Eqs.(2) and (9) we

immediately obtain

$$\begin{aligned}
& \frac{1}{2} \Delta^{[s]T} Q^{[s]} \Delta^{[s]} + \Delta^{[s]T} C^{[s]} \\
&= \gamma \sum_{r=1}^{n-1} \left(\frac{f_{r+1}^{[s-1]} - f_r^{[s-1]}}{\xi_r} \right) \left(\frac{\Delta_{r+1}^{[s]} - \Delta_r^{[s]}}{\xi_r} \right) \xi_r + \frac{\gamma}{2} \sum_{r=1}^{n-1} \left(\frac{\Delta_{r+1}^{[s]} - \Delta_r^{[s]}}{\xi_r} \right)^2 \xi_r \\
&+ 4\varphi \sum_{r=2}^{n-1} \left(\frac{1}{(\xi_{r-1} + \xi_r)} \left(\frac{1}{\xi_r} f_{r+1}^{[s-1]} + \frac{1}{\xi_{r-1}} f_{r-1}^{[s-1]} \right) - \frac{1}{\xi_r \xi_{r-1}} f_r^{[s-1]} \right) \\
&\times \left(\frac{1}{(\xi_{r-1} + \xi_r)} \left(\frac{1}{\xi_r} \Delta_{r+1}^{[s]} + \frac{1}{\xi_{r-1}} \Delta_{r-1}^{[s]} \right) - \frac{1}{\xi_r \xi_{r-1}} \Delta_r^{[s]} \right) \xi_r \\
&+ 2\varphi \sum_{r=2}^{n-1} \left(\frac{1}{(\xi_{r-1} + \xi_r)} \left(\frac{1}{\xi_r} \Delta_{r+1}^{[s]} + \frac{1}{\xi_{r-1}} \Delta_{r-1}^{[s]} \right) - \frac{1}{\xi_r \xi_{r-1}} \Delta_r^{[s]} \right)^2 \xi_r,
\end{aligned}$$

and defining

$$Y_\lambda^{[s]} := \sum_{j=1}^m \lambda_j^{[s]} \left[\hat{\rho}_j^{[s]} - \ln(v_j^{[s]}) + \sum_{r=1}^{R_{n_j}^{(j)}-1} \hat{f}_r^{[s]} \xi_r \right], \quad (10)$$

$$Y_\mu^{[s]} := -\mu^{[s-1]} \sum_{r=1}^n \ln(\hat{f}_r^{[s]}), \quad (11)$$

$$Y_{\tilde{\mu}}^{[s]} := -\tilde{\mu}^{[s-1]} \sum_{j=1}^m \left(\ln(\rho_j^{[s]} - \rho_j^b) + \ln(\rho_j^a - \rho_j^{[s]}) \right), \quad (12)$$

we have

$$\begin{aligned}
Y_\lambda^{[s]} &= \sum_{j=1}^m \lambda_j^{[s]} \left[\rho_j^{[s-1]} + \sigma_j^{[s]} - \ln(v_j^{[s-1]}) + \sum_{r=1}^{R_{n_j}^{(j)}-1} (f_r^{[s-1]} + \Delta_r^{[s]}) \xi_r \right. \\
&\quad \left. - \frac{1}{v_j^{[s-1]}} \sum_{i=1}^{n_j-1} \sum_{r=R_i^{(j)}}^{R_{n_j}^{(j)}-1} \alpha_{ij} \exp \left(\sum_{z=R_i^{(j)}}^{R_{n_j}^{(j)}-1} f_z^{[s-1]} \xi_z \right) \Delta_r^{[s]} \xi_r + O(\Delta_r^{[s]2}) \right], \\
Y_\mu^{[s]} &= -\mu^{[s-1]} \sum_{r=1}^n \left(\frac{\Delta_r^{[s]}}{f_r^{[s-1]}} - \frac{1}{2} \left(\frac{\Delta_r^{[s]}}{f_r^{[s-1]}} \right)^2 + O(\Delta_r^{[s]3}) \right), \\
Y_{\tilde{\mu}}^{[s]} &= -\tilde{\mu}^{[s-1]} \sum_{j=1}^m \left(\ln(\rho_j^{[s-1]} - \rho_j^b) + \ln(\rho_j^a - \rho_j^{[s-1]}) \right) \\
&\quad + \tilde{\mu}^{[s-1]} \sum_{j=1}^m \left(\frac{1}{\rho_j^a - \rho_j^{[s-1]}} - \frac{1}{\rho_j^{[s-1]} - \rho_j^b} \right) \sigma_j^{[s]} \\
&\quad + \frac{\tilde{\mu}^{[s-1]}}{2} \sum_{j=1}^m \left[\frac{1}{(\rho_j^{[s-1]} - \rho_j^a)^2} + \frac{1}{(\rho_j^{[s-1]} - \rho_j^b)^2} \right] \sigma_j^{[s]2} + O(\sigma_j^{[s]3}).
\end{aligned}$$

Thus defining

$$\chi(r, x, y) := \begin{cases} 1 & x \leq r \leq y \\ 0 & \text{otherwise} \end{cases}, \quad \delta_{i,j} := \begin{cases} 1 & i = j \\ 0 & \text{otherwise} \end{cases},$$

from above expansions and Eq.(9) we immediately obtain

$$\begin{aligned} B_{r,j}^{[s]} &= \xi_r \chi(r, 1, R_{n_j}^{(j)} - 1) - \frac{1}{v_j^{[s-1]}} \sum_{i=1}^{n_j-1} \alpha_{ij} \exp \left(\sum_{z=R_i^{(j)}}^{R_{n_j}^{(j)}-1} f_z^{[s-1]} \xi_z \right) \xi_r \chi(r, R_i^{(j)}, R_{n_j}^{(j)} - 1), \\ a_j^{[s]} &= \rho_j^{[s-1]} - \ln(v_j^{[s-1]}) + \sum_{r=1}^{R_{n_j}^{(j)}-1} f_r^{[s-1]} \xi_r, \\ Q_{r,x}^{[s]} &= 4\varphi \left[\left(\frac{(1-\delta_{r,1})(1-\delta_{r,2})\xi_{r-1}}{(\xi_{r-2} + \xi_{r-1})^2 \xi_{r-1}^2} + \frac{(1-\delta_{r,n})(1-\delta_{r,1})\xi_r}{\xi_r^2 \xi_{r-1}^2} + \frac{(1-\delta_{r,n-1})(1-\delta_{r,n})\xi_{r+1}}{(\xi_r + \xi_{r+1})^2 \xi_r^2} \right) \delta_{r,x} \right. \\ &\quad + \frac{\xi_{r+1}\delta_{r+2,x}}{(\xi_r + \xi_{r+1})^2 \xi_{r+1}\xi_r} + \frac{\xi_{x+1}\delta_{r-2,x}}{(\xi_x + \xi_{x+1})^2 \xi_{x+1}\xi_x} - \frac{(1-\delta_{r,1})\xi_r\delta_{r+1,x}}{(\xi_{r-1} + \xi_r)\xi_{r-1}\xi_r^2} \\ &\quad - \frac{(1-\delta_{x,1})\xi_x\delta_{r-1,x}}{(\xi_{x-1} + \xi_x)\xi_{x-1}\xi_x^2} - \frac{(1-\delta_{r,n})\xi_r\delta_{r-1,x}}{(\xi_{r-1} + \xi_r)\xi_r\xi_{r-1}^2} - \frac{(1-\delta_{x,n})\xi_x\delta_{r+1,x}}{(\xi_{x-1} + \xi_x)\xi_x\xi_{x-1}^2} \Big] \\ &\quad + \gamma \left[(1-\delta_{r,n})\frac{\xi_r}{\xi_r^2}\delta_{r,x} + (1-\delta_{r,1})\frac{\xi_{r-1}}{\xi_{r-1}^2}\delta_{r,x} - \frac{\xi_r}{\xi_r^2}\delta_{r+1,x} - \frac{\xi_x}{\xi_x^2}\delta_{r-1,x} \right] + \frac{\mu^{[s-1]}}{f_r^{[s-1]2}}\delta_{r,x}, \\ C_r^{[s]} &= -\frac{\mu^{[s-1]}}{f_r^{[s-1]}} + \gamma \frac{f_r^{[s-1]} - f_{r-1}^{[s-1]}}{\xi_{r-1}} \frac{(1-\delta_{r,1})}{\xi_{r-1}} \xi_{r-1} - \gamma \frac{f_{r+1}^{[s-1]} - f_r^{[s-1]}}{\xi_r} \frac{(1-\delta_{r,n})}{\xi_r} \xi_r \\ &\quad + 4\varphi \left(\frac{f_r^{[s-1]} - f_{r-1}^{[s-1]}}{\xi_{r-1}} + \frac{f_{r-2}^{[s-1]} - f_{r-1}^{[s-1]}}{\xi_{r-2}} \right) \frac{(1-\delta_{r,1})(1-\delta_{r,2})}{(\xi_{r-2} + \xi_{r-1})^2 \xi_{r-1}} \xi_{r-1} \\ &\quad + 4\varphi \left(\frac{f_{r+2}^{[s-1]} - f_{r+1}^{[s-1]}}{\xi_{r+1}} + \frac{f_r^{[s-1]} - f_{r+1}^{[s-1]}}{\xi_r} \right) \frac{(1-\delta_{r,n})(1-\delta_{r,n-1})}{(\xi_r + \xi_{r+1})^2 \xi_r} \xi_{r+1} \\ &\quad - 4\varphi \left(\frac{f_{r+1}^{[s-1]} - f_r^{[s-1]}}{\xi_r} + \frac{f_{r-1}^{[s-1]} - f_r^{[s-1]}}{\xi_{r-1}} \right) \frac{(1-\delta_{r,1})(1-\delta_{r,n})}{(\xi_{r-1} + \xi_r)\xi_r\xi_{r-1}} \xi_r, \\ M_{j,k}^{[s]} &= \tilde{\mu}^{[s-1]} \left[\frac{1}{(\rho_j^a - \rho_j^{[s-1]})^2} + \frac{1}{(\rho_j^b - \rho_j^{[s-1]})^2} \right] \delta_{j,k}, \\ D_j^{[s]} &= \tilde{\mu}^{[s-1]} \left(\frac{1}{\rho_j^a - \rho_j^{[s-1]}} + \frac{1}{\rho_j^b - \rho_j^{[s-1]}} \right). \end{aligned}$$

Appendix B. Dynamic Programming for a quadratic objective

Given positive definite symmetric matrices $Q \in R^{n,n}$ and $M \in R^{c,c}$ and the n -vector C and c -vector D , we want to minimize the objective function

$$W(\Delta, \sigma) := \frac{1}{2} \Delta^T Q \Delta + \Delta^T C + \frac{1}{2} \sigma^T M \sigma + \sigma^T D,$$

subject to the set of constraints

$$a_j + \sigma_j + \sum_{k=0}^n \Delta_k B_{k,j} = 0, \quad j = 1, \dots, c, \quad (13)$$

thus defining a new objective function

$$Z(\Delta, \sigma, \lambda) := W(\Delta, \sigma) + (\Delta^T B + a^T + \sigma^T) \lambda.$$

For any Q we define

$$d := \max_k \left(\sum_{s=1}^{k-1} \delta(Q_{k,s}) \right), \quad \delta(x) := \begin{cases} 1 & x \neq 0 \\ 0 & x = 0 \end{cases},$$

where e.g. $d = 0, 1$ for diagonal, tridiagonal matrices respectively. When $d \ll n$ and $c \ll n$ then Z can be solved efficiently with dynamic programming. To set up the notation and for those readers not familiar with dynamic programming let us here briefly explain the basics of this well known method [19]. We start defining

$$\begin{aligned} Z^{(n)} &:= Z, \\ Z^{(q-1)} &:= Z^{(q)} \Big|_{\Delta_q = \Delta_q^*}, \quad q = 1, \dots, n, \end{aligned} \quad (14)$$

where Δ_q^* satisfies

$$\nabla_{\Delta_q} Z^{(q)} \Big|_{\Delta_q = \Delta_q^*} = 0, \quad q = 1, \dots, n. \quad (15)$$

For $q = 0, \dots, n$ we use the inductive hypothesis

$$\begin{aligned} Z^{(q)} &= \frac{1}{2} \Delta^T Q^{(q)} \Delta + \Delta^T B^{(q)} \lambda + \Delta^T C^{(q)} + \lambda^T a^{(q)} \\ &\quad + \frac{1}{2} \lambda^T G^{(q)} \lambda + b^{(q)} + \sigma^T \lambda + \frac{1}{2} \sigma^T M \sigma + \sigma^T D, \end{aligned} \quad (16)$$

where $Q^{(q)} \in R^{q,q}$, $B^{(q)} \in R^{c,q}$ and $G^{(q)} \in R^{c,c}$. The final step of this backwards process consists in obtaining σ^* and λ^* satisfying

$$\begin{aligned} \nabla_{\sigma} Z^{(0)} \Big|_{\sigma = \sigma^*} &= 0, \\ \nabla_{\lambda} Z^{(0)} \Big|_{\lambda = \lambda^*} &= 0. \end{aligned} \quad (17)$$

From Eqs.(15) and (16) we obtain

$$\Delta_q^* = -\frac{1}{Q_{q,q}^{(q)}} \left(\sum_{r=q-d}^{q-1} Q_{q,r}^{(q)} \Delta_r + B_q^{(q)} \lambda + C_q^{(q)} \right), \quad (18)$$

and plugging Eq.(18) into Eq.(14) we obtain

$$\begin{aligned}
Z^{(q-1)} &= Z^{(q)} \Big|_{\Delta_q=0} - \frac{1}{2} \frac{1}{Q_{q,q}^{(q)}} \left(\sum_{r=q-d}^{q-1} Q_{q,r}^{(q)} \Delta_r + B_q^{(q)} \lambda + C_q^{(q)} \right)^2 \\
&= \frac{1}{2} \sum_{\{r,s\}=1}^{q-1} \Delta_r \left(Q_{r,s}^{(q)} - \frac{Q_{q,r}^{(q)} Q_{q,s}^{(q)}}{Q_{q,q}^{(q)}} \theta(r-q+d) \theta(s-q+d) \right) \Delta_s \\
&\quad + \sum_{r=1}^{q-1} \Delta_r \left(\left(B_r^{(q)} - \frac{Q_{q,r}^{(q)} B_q^{(q)}}{Q_{q,q}^{(q)}} \theta(r-q+d) \right) \lambda + C_r^{(q)} - \frac{Q_{q,r}^{(q)} C_q^{(q)}}{Q_{q,q}^{(q)}} \theta(r-q+d) \right) \\
&\quad + \frac{1}{2} \lambda^T \left(G^{(q)} - \frac{B_q^{(q)T} B_q^{(q)}}{Q_{q,q}^{(q)}} \right) \lambda + \left(\alpha^{(q)T} - \frac{C_q^{(q)} B_q^{(q)}}{Q_{q,q}^{(q)}} \right) \lambda - \frac{1}{2} \frac{C_q^{(q)2}}{Q_{q,q}^{(q)}} + b^{(q)} \\
&\quad + \sigma^T \lambda + \frac{1}{2} \sigma^T M \sigma + \sigma^T D,
\end{aligned}$$

hence obtaining

$$Q_{r,s}^{(q-1)} = Q_{r,s}^{(q)} - \frac{Q_{q,r}^{(q)} Q_{q,s}^{(q)}}{Q_{q,q}^{(q)}} \theta(r-q+d) \theta(s-q+d), \quad (19)$$

$$B_r^{(q-1)} = B_r^{(q)} - \frac{Q_{q,r}^{(q)} B_q^{(q)}}{Q_{q,q}^{(q)}} \theta(r-q+d), \quad (20)$$

$$C_r^{(q-1)} = C_r^{(q)} - \frac{Q_{q,r}^{(q)} C_q^{(q)}}{Q_{q,q}^{(q)}} \theta(r-q+d), \quad (21)$$

$$G^{(q-1)} = G^{(q)} - \frac{B_q^{(q)T} B_q^{(q)}}{Q_{q,q}^{(q)}}, \quad (22)$$

$$a^{(q-1)} = a^{(q)} - \frac{C_q^{(q)} B_q^{(q)}}{Q_{q,q}^{(q)}}, \quad (23)$$

$$b^{(q-1)} = b^{(q)} - \frac{1}{2} \frac{C_q^{(q)2}}{Q_{q,q}^{(q)}}, \quad (24)$$

where

$$\theta(x) := \begin{cases} 1 & x \geq 0, \\ 0 & x < 0. \end{cases}$$

Now from Eq.(17) we obtain

$$\begin{aligned}
M\sigma^* + \lambda^* + D &= 0, \\
G^{(0)}\lambda^* + \sigma^* + a^{(0)} &= 0,
\end{aligned}$$

or

$$\begin{aligned}\lambda^* &= \left(M^{-1} - G^{(0)}\right)^{-1} \left(a^{(0)} - M^{-1}D\right) = \left(I - MG^{(0)}\right)^{-1} \left(Ma^{(0)} - D\right), \\ \sigma^* &= -M^{-1}(\lambda^* + D),\end{aligned}\tag{25}$$

and using Eq.(22) we obtain

$$\lambda^* = \left(M^{-1} + \sum_{q=1}^n \frac{B_q^{(q)T} B_q^{(q)}}{Q_{q,q}^{(q)}}\right)^{-1} \left(a^{(0)} - M^{-1}D\right) = \left(I + M \sum_{q=1}^n \frac{B_q^{(q)T} B_q^{(q)}}{Q_{q,q}^{(q)}}\right)^{-1} \left(Ma^{(0)} - D\right).$$

Finally from λ^* and Eq.(25) we obtain σ^* and then using Eqs.(18) and (19-21) we obtain Δ^* moving forward in the q index.

B.1 Computing times of a dynamic programming iteration

The time necessary to compute all $Q^{(i)}$ is proportional to

$$T_1 = (n-d) \frac{d(d+1)}{2} + \sum_{i=1}^{d-1} \frac{i(i+1)}{2} = \frac{d(d+1)}{2} \left(n - \frac{2d+1}{3}\right).$$

To calculate the inverse of $M^{-1} - G^{(0)}$, that is a $c \times c$ symmetric matrix, we require a computing time proportional to

$$T_2 = \frac{1}{6} (c^3 - c).$$

All $C^{(i)}$, $B^{(i)}$ and $a^{(i)}$ require, respectively, computing times proportional to

$$T_3 = (n-d)d + \sum_{i=1}^{d-1} i = \frac{d}{2} (2n-d-1),$$

$$T_4 = cT_3,$$

$$T_5 = cn.$$

Finally the calculation of $G^{(0)}$ requires a computing time proportional to

$$T_6 = \frac{c(c+1)}{2}n.$$

In the forward rate calculation typically we have $n > c > d$ (in Eq.(2) we have $d = 2$) and therefore the maximum delay would be given by T_6 .

Appendix C. Fulfilling inequalities and updating log-barriers

The Newton step obtained from Eq.(8) may not fulfill the inequality constraints (4). To satisfy such constraints we search for a $\alpha^{[s]}$ in the interval $[0, 1]$ such that

$$f_r^{[s-1]} + \alpha^{[s]} \Delta_r^{[s]} \geq 0, \quad \rho_j^b \leq \rho_j^{[s-1]} + \alpha^{[s]} \sigma_j^{[s]} \leq \rho_j^a.\tag{26}$$

In order to do this we first determine the maximum $\alpha_{\max}^{[s]}$ in the interval $[0, 1]$ satisfying Eq.(26). That is, given the sets of points

$$\begin{aligned} A &:= \left\{ r, \hat{f}_r^{[s]} \leq 0 \right\}, \\ \tilde{A}_{\leq} &:= \left\{ j, \hat{\rho}_j^{[s]} \leq \rho_j^b \right\}, \\ \tilde{A}_{\geq} &:= \left\{ j, \hat{\rho}_j^{[s]} \geq \rho_j^a \right\}, \end{aligned}$$

we have

$$\alpha_{\max}^{[s]} = \min \left(\min_{r \in A} \left(\frac{-f_r^{[s-1]}}{\Delta_r^{[s]}} \right), \min_{j \in \tilde{A}_{\leq}} \left(\frac{\rho_j^b - \rho_j^{[s-1]}}{\sigma_j^{[s]}} \right), \min_{j \in \tilde{A}_{\geq}} \left(\frac{\rho_j^a - \rho_j^{[s-1]}}{\sigma_j^{[s]}} \right) \right),$$

and then we take

$$\alpha^{[s]} := \beta \alpha_{\max}^{[s]},$$

with $0 < \beta < 1$ (in our implementation we have taken $\beta = .9$ that is a standard election in the optimization literature). Once we have $\alpha^{[s]}$ we define

$$\begin{aligned} f_r^{[s]} &= f_r^{[s-1]} + \alpha^{[s]} \Delta_r^{[s]}, \\ \rho_j^{[s]} &= \rho_j^{[s-1]} + \alpha^{[s]} \sigma_j^{[s]}, \\ \mu^{[s]} &= \max(\Psi(\alpha^{[s]}) \mu^{[s-1]}, \mu_{\min}), \\ \tilde{\mu}^{[s]} &= \max(\Psi(\alpha^{[s]}) \tilde{\mu}^{[s-1]}, \tilde{\mu}_{\min}), \end{aligned} \tag{27}$$

where $\mu_{\min}$ and $\tilde{\mu}_{\min}$ are positive small values guaranteeing that matrices Q and M are positive definite and $\Psi : [0, 1] \rightarrow R^+$ is a monotonically decreasing function satisfying $\Psi(0) = 1$. In our implementation we have taken Ψ of the form

$$\Psi(\alpha) = (1 - l)(1 - \alpha)^{\xi} + l, \tag{28}$$

with $l = 10^{-2}$, $\xi = 1$. The value of l controls how fast barriers are reduced. In our implementation we have found that adequate values for l range $10^{-3} \lesssim l \lesssim 10^{-1}$. The value of ξ controls the non-linearity of Ψ . We have found that the simple linear response provides good performance albeit convergence time is not significantly affected for ξ in the range $0.6 \lesssim \xi \lesssim 2$

In this way we iterate the algorithm n_{it} times until a given termination criterion is met. Defining

$$\begin{aligned} \delta_{\ln W}^{[s]} &:= \ln \left(W \left(f^{[s]} \right) \right) - \ln \left(W \left(f^{[s-1]} \right) \right), \\ \epsilon^{[s]} &:= \max_j \left(\left| \rho_j^{[s]} - \ln \left(v_j^{[s]} \right) + \sum_{r=1}^{R_{n_j}^{(j)}-1} f_r^{[s]} \xi_r \right| \right), \end{aligned}$$

we have chosen the following termination criterion

$$n_{it} = N_{\max} \text{ or } \left\{ \left[W^{[s]} < W_{zero} \text{ or } \left(\left| \delta_{\ln W}^{[s]} \right| < \delta_{\ln W}^{\max} \text{ and } \left| \delta_{\ln W}^{[s-1]} \right| < \delta_{\ln W}^{\max} \right) \right] \right. \\ \left. \text{and } n_{it} > N_{\min} \text{ and } \mu^{[s]} < \mu_{\max} \text{ and } \tilde{\mu}^{[s]} < \tilde{\mu}_{\max} \text{ and } \epsilon^{[s]} < \epsilon_{\max} \right\}. \quad (29)$$

In our implementation we have taken

$$\begin{aligned} \mu^{[0]} &= 10^{-1}, & \mu_{\min} &= 10^{-10}, & \mu_{\max} &= 10^{-6}, \\ \tilde{\mu}^{[0]} &= 10^{+1}, & \tilde{\mu}_{\min} &= 10^{-10}, & \tilde{\mu}_{\max} &= 10^{-6}, \\ \epsilon_{\max} &= 10^{-8}, & \delta_{\ln W}^{\max} &= 10^{-2}, & W_{zero} &= 10^{-9}, \\ N_{\min} &= 5, & N_{\max} &= 60. \end{aligned} \quad (30)$$

Obviously there is considerable latitude to change the heuristic values assigned to the above parameters. Let us finish this appendix making some comments regarding their robustness.

$N_{\min}$ is there to guarantee a minimum number of iterations so as to have $\delta_{\ln W}^{[s]}$ and $\delta_{\ln W}^{[s-1]}$ well defined and to avoid premature termination in the improbable case where the other criteria incorrectly suggest convergence. For this purpose is enough to take $2 \leq N_{\min} \lesssim 5$.

$N_{\max}$ serves as a maximum limit to secure termination even if convergence is not achieved and therefore an alarm should be provided whenever $n_{it} = N_{\max}$. From our experience we observe that is more than enough to take $50 \lesssim N_{\max} \lesssim 100$.

W_{zero} sets our precision to consider a given forward rate curve as a straight line. The order of magnitude of W_{zero} should be taken much lower than the typical order of magnitude of the observed optimal W . The value of the optimal W depends not only on the constraining data but also on γ and φ . We have adopted the practise of spanning the range $0 \leq \sqrt{\gamma/\varphi} \leq 1 \text{ year}^{-1}$ taking $0 \leq \gamma \leq 1 \text{ year}^3$, $\varphi = 1 \text{ year}^5$ and the range $1 \text{ year}^{-1} \leq \sqrt{\gamma/\varphi} \leq \infty$ taking $\gamma = 1 \text{ year}^3$, $0 \leq \varphi \leq 1 \text{ year}^5$. With this convention we have found reasonable to take $10^{-10} \lesssim W_{zero} \lesssim 10^{-8}$ for the full range $0 \leq \sqrt{\gamma/\varphi} \leq \infty$.

$\delta_{\ln W}^{\max}$ sets the maximum variation of $\ln(W)$ between newton steps that is accepted before termination. Note that in (29) to have $\left(\left| \delta_{\ln W}^{[s]} \right| < \delta_{\ln W}^{\max} \text{ and } \left| \delta_{\ln W}^{[s-1]} \right| < \delta_{\ln W}^{\max} \right)$ is a necessary but not sufficient condition for termination. Therefore sending $\delta_{\ln W}^{\max} \rightarrow \infty$ has no major impact and is equivalent to rely only on the error $\epsilon^{[s]}$ and the barrier coefficients as indicators of convergence. On the contrary excessively reducing $\delta_{\ln W}^{\max}$ can generate unnecessary iterations. We have tested that for values satisfying $\delta_{\ln W}^{\max} \gtrsim 10^{-5}$ we do not have any drastic increase in convergence time.

$\epsilon_{\max}$ controls the error in the constraints and is the most important parameter in (29). A too large value of $\epsilon_{\max}$ reduces the accuracy of the result and a too small value can give

rise to unnecessary iterations. We have observed acceptable results for $\epsilon_{\max}$ in the range $10^{-4} \lesssim \epsilon_{\max} \lesssim 10^{-10}$.

$\mu_{\max}$ and $\tilde{\mu}_{\max}$ control the maximum allowed values for the logarithmic barriers implementing the positivity and spread constraints respectively. Large values for these parameters can make log barriers to have a residual influence in the feasible region. Acceptable values of these maximum weights range in $\mu_{\min} < \mu_{\max} \lesssim 10^{-6}$ and $\tilde{\mu}_{\min} < \tilde{\mu}_{\max} \lesssim 10^{-6}$. As explained above keeping parameters $\mu_{\min}$ and $\tilde{\mu}_{\min}$ positive guarantees that matrices Q and M are positive definite which in turn is sufficient to guarantee the existence of an optimal solution in each iteration. Hence $\mu_{\min}$ and $\tilde{\mu}_{\min}$ should be chosen as small as possible without interfering with the numerical stability of the algorithm. Using double precision in our program we have found the values given in (29) as a good compromise.

Finally $\mu^{[0]}$ and $\tilde{\mu}^{[0]}$ are the initial values for the log barriers coefficients. Taking very large values for $\mu^{[0]}$ and $\tilde{\mu}^{[0]}$ increases the convergence time because we need more time to reduce the barriers. Taking too small values for $\mu^{[0]}$ and $\tilde{\mu}^{[0]}$ also increases the convergence time because time is wasted exploring unfeasible solutions. Moreover, we have observed that convergence and stability is improved if the contributions to the objective of the two barrier terms are kept balanced. This is achieved setting $\tilde{\mu}^{[0]} \simeq \frac{n}{2m}\mu^{[0]}$ (see Eqs.(11) and (12)) and using the same updating factor $\Psi(\alpha^{[s]})$ for $\mu^{[s]}$ and $\tilde{\mu}^{[s]}$ (see Eq.(27)). Keeping $\tilde{\mu}^{[0]} \simeq \frac{n}{2m}\mu^{[0]}$ we have found that the time of convergence is stable for $\mu^{[0]}$ in the range $10^{-4} \lesssim \mu^{[0]} \lesssim 10^{+1}$.

Appendix D. Bond tables

In this work we have used the following data tables and conventions.

The price of the bond j is calculated using the formula

$$P_j = \sum_{i=1}^{n_j} \frac{(c_j/100 + \delta_{i,n_j}) N_j}{(1 + \frac{r_j}{100})^{\Delta T_i}}, \quad (31)$$

where N_j is the nominal amount (SEK 40 millions for all bonds in Table 1), c_j is the coupon rate (given in Table 2), n_j is the total number of remaining coupons (each paid at time T_i^j), r_j is the quoted rate given in Table 1 and ΔT_i is a time difference between T_i^j and the settlement day. This time difference is calculated according to the ISMA 30E/360 convention defined as follows: given two dates (d_1, m_1, y_1) and (d_2, m_2, y_2) , their ISMA 30E/360 time difference ΔT is given by

$$\begin{aligned} & \text{if } d_1 = 31 \text{ set } d_1 \text{ to } 30, \\ & \text{if } d_2 = 31 \text{ set } d_2 \text{ to } 30, \\ & \Delta T = y_2 - y_1 + \frac{30(m_2 - m_1) + d_2 - d_1}{360}, \end{aligned}$$

Date \ Bond (SO)	1033	1042	1035	1044	1038	1037	1040	1043	1034	1045	1041
Friday 06	4.86	4.92	5.06	5.15	5.26	5.27	5.355	5.4	5.395	5.46	5.655
Monday 09	4.905	4.965	5.11	5.2	5.05	5.325	5.4	5.455	5.23	5.51	5.7
Tuesday 10	4.885	4.945	5.095	5.185	4.93	5.295	5.395	5.435	5.24	5.495	5.68
Wednesday 11	4.865	4.92	5.06	5.15	4.93	5.275	5.355	5.41	5.415	5.465	5.65
Thursday 12	4.835	4.885	5.025	5.125	4.93	5.25	5.355	5.405	5.405	5.465	5.66
Friday 13	4.84	4.89	5.045	5.145	4.93	5.27	5.38	5.43	5.43	5.495	5.69
Monday 16	4.825	4.87	5.035	5.13	4.93	5.25	5.36	5.415	5.41	5.475	5.66
Tuesday 17	4.805	4.85	5.015	5.11	4.93	5.23	5.34	5.395	5.395	5.455	5.64
Wednesday 18	4.8	4.86	5.005	5.1	4.93	5.22	5.335	5.38	5.395	5.445	5.64
Thursday 19	4.77	4.83	4.97	5.06	4.93	5.165	5.27	5.34	5.34	5.39	5.585

Table 1: Quoted rates for the eleven Swedish Government Bonds used in the calculation of the forward rate curves corresponding to Figs.(1-3). All days are from a two-week period of July 2001. The description of each of the bonds is given in Table 2. The prices of the bonds are obtained from this rates and the data of Table 2 using Eq.(31).

References

- [1] McCulloch, J. H., *Measuring the term structure of interest rates*, Journal of Business, 44, 19-31, 1971. McCulloch, J. H., *The Tax-Adjusted Yield Curve*, Journal of Finance, 30, 811-830, 1975.
- [2] Vasicek, O. A. and Fong, H. G. *Term structure modeling using exponential splines*, Journal of Finance, 37(2), 329-348, 1982.
- [3] Chambers, D., Carleton, W. and Waldman, D., *A New Approach to Estimation of the Term Structure of Interest Rates*, Journal of Financial and Quantitative Analysis 19(3), 223-269, 1984.
- [4] Shea, G., *Interest Rate Term Structure Estimation with Exponential Splines: A Note*, Journal of Finance, 40, 319-325, 1985.
- [5] Nelson, C. and Siegel, A., *Parsimonious Modeling of Yield Curves*, Journal of Business 60, 473-489, 1987.

Bond	Maturity (dd/mm/yyyy)	annual coupon (%)
SO 1033	05/05/2003	10.25
SO 1042	15/01/2004	5
SO 1035	09/02/2005	6
SO 1044	20/04/2006	3.5
SO 1038	25/10/2006	6.5
SO 1037	15/08/2007	8
SO 1040	05/05/2008	6.5
SO 1043	28/01/2009	5
SO 1034	20/04/2009	9
SO 1045	15/03/2011	5.25
SO 1041	05/05/2014	6.75

Table 2: Maturities and coupon rates for the 11 Swedish bonds quoted in Table 1. Coupons are paid annually with the last coupon paid at maturity together with a nominal of SEK 40 millions.

- [6] Svensson, L. O., *Estimating and Interpreting Forward Interest Rates: Sweden 1992-1994*. Technical Report Discussion Paper Number 1051, Centre for Economic Policy Research, October 1994.
- [7] Fisher, M., Nychka D. and Zervos D., *Fitting the term structure of interest rates with smoothing splines*, FEDS 95-1, Federal Reserve Board, Washington DC. 1994.
- [8] Waggoner D. F., *Spline Methods for Extracting Interest Rates Curves from Coupon Bond Prices*, Working Paper 97-10, Federal Reserve Bank of Atlanta, November 1997.
- [9] Adams, K. and van Deventer, D., *Fitting Yield Curves and Smooth Forward Rate Curves with Maximum Smoothness*, Journal of Fixed Income, V4, 53-62, June 1994.
- [10] Delbaen F., Lorimier S. *Estimation of the yield curve and the forward rate curve starting from a finite number of observations*. - In: Insurance: mathematics and economics, 11:4, 249-258 1992.
- [11] Kian Guan Lim and Qin Xiao, *Computing Maximum Smoothness Forward Rate Curves*, Statistics and Computing, Kluwer Academic Publishers, Vol.12, pp. 275-279, ISSN 0960-3174, 2002.

- [12] Kian Guan Lim, Qin Xiao, and Jimmy Ang, *Estimating Forward Rate Curve in Pricing Interest Rate Derivatives*, Derivatives Use, Trading & Regulation, an International Journal of the Futures and Options Association UK, Vol. 6 No.4, pp. 299-305, 2001.
- [13] Frishling V. and Yamamura, J., *Fitting a Smooth Forward Rate Curve to Coupon Instruments*, Journal of Fixed Income, V6 12, 97-103, 1996.
- [14] Yekutieli, I., *With bond Stripping, the Curve's the Thing*, Bloomberg Magazine 8(3), 84-90, 1999.
- [15] Forsgren A., *A note on maximum-smoothness approximation of forward interest rate*, Report TRITA-MAT-1998-OS3, Department of Mathematics, Royal Institute of Technology, 1998.
- [16] Kwon, Oh Kang, *A general framework for the construction and the smoothing of forward rate curves*. QFRG, University of Technology, Sydney. <http://www.business.uts.edu.au/finance/qfr/cfrg-papers.html>, March 2002.
- [17] Wets, Roger J. B., Bianchi, Stephen W. and Yang, Liming, *Serious Zero-Curves*, EpiSolutions Inc., <http://www.episolutionsinc.com/html/downloads/SeriousZC.pdf>, January 2002.
- [18] see for example: J. Nocedal and S. J. Wright. *Numerical Optimization*. Springer Verlag, 1999.
- [19] see for example: Bertsekas, Dimitri P., *Dynamic Programming and Optimal Control: Vol.1*, 2nd Edition, Athena Scientific, 2000.
- [20] Weiss, S.M. and Kulikowski, C.A., *Computer Systems That Learn*, Morgan Kaufmann, 1991.
- [21] Stone, M., *Asymptotics for and against cross-validation*, Biometrika, 64, 29-35, 1977.